



Principal components analysis



Herman Kamper



2024-02, CC BY-SA 4.0

Basic idea







View 1: Maximising variance





View 2: Minimising reconstruction error





Preliminaries

Lagrange multipliers

Want to optimise $f(x)$ subject to some constraint $g(x) = 0$.

We define a new objective:

$$J(x, \lambda) = f(x) + \lambda g(x)$$

and optimise w.r.t. both x and λ .

Eigenvalues and eigenvectors

For a square matrix \mathbf{A} :

$$\mathbf{A}\mathbf{u} = \lambda\mathbf{u}$$

The solutions to this equation are pairs of eigenvalues λ with eigenvectors \mathbf{u} .

Vector derivatives

See the notes and videos on vector and matrix derivatives (03b).

PCA setup

We want to project $\mathbf{x}^{(n)} \in \mathbb{R}^D$ to $\mathbf{z}^{(n)} \in \mathbb{R}^M$, with $M < D$.

(We normally assume that the inputs \mathbf{x} have been normalised to have a zero mean.)



To do the projection, we use M projection vectors $\mathbf{w}_m \in \mathbb{R}^D$.

The projection vectors $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_M$ are unit length and orthogonal:

$$\|\mathbf{w}_m\| = 1 \quad \text{and} \quad \mathbf{w}_i^\top \mathbf{w}_j = 0 \quad \forall i \neq j$$

The projection of the n^{th} item $\mathbf{x}^{(n)}$ onto the m^{th} dimension is

$$z_m^{(n)} = \mathbf{w}_m^\top \mathbf{x}^{(n)}$$




Projection

So $\mathbf{x}^{(n)}$ is mapped to

$$\mathbf{z}^{(n)} =$$

Reconstruction



Finding the projection vectors


Setup

- Data that have been mean-normalised: $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots, \mathbf{x}^{(N)}$
- Want to find projection vectors: $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_M$
- Unit length: $\|\mathbf{w}_m\| = 1$
- Orthogonal: $\mathbf{w}_i^\top \mathbf{w}_j = 0 \quad \forall i \neq j$

Objective

Want to find $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_M$ so that (sample) variance is maximised.

Let's first just look at one dimension.



$$\hat{\sigma}_{z_1}^2 =$$

We want to maximise $\hat{\sigma}_{z_1}^2$ subject to $\|\mathbf{w}_1\| = 1$, i.e. $\mathbf{w}_1^\top \mathbf{w}_1 = 1$. We use a Lagrange multiplier:

$$J(\mathbf{w}_1) =$$

Minimise the loss w.r.t. \mathbf{w}_1 :

Which eigenvector-value do we use?

So pick eigenvector corresponding to the largest eigenvalue.


How do we find \mathbf{w}_2 , with $\|\mathbf{w}_2\| = 1$, and $\mathbf{w}_1^\top \mathbf{w}_2 = 0$? Repeat above steps:

$$\hat{\Sigma} \mathbf{w}_2 = \lambda_2 \mathbf{w}_2$$

So pick eigenvector corresponding to the second largest eigenvalue.
Etc.

PCA view 2: Minimising the reconstruction error

Instead of maximising variance, we think of PCA as minimising the reconstruction loss:



Let's show that these two views are the same:

$$J(\mathbf{w}_1) =$$


(Here we looked at projection to one dimension, but you can follow the same steps for projection to $M > 1$ dimensions.)

The relationship of PCA to SVD

(Not examinable.)

Singular value decomposition (SVD)

$$\mathbf{X} = \mathbf{U}\mathbf{S}\mathbf{V}^T$$



Relationship to PCA


Take SVD of the design matrix \mathbf{X} :

$$\mathbf{X} = \mathbf{U}\mathbf{S}\mathbf{V}^T$$

Then

$$\begin{aligned} \mathbf{X}^T \mathbf{X} &= \mathbf{V} \mathbf{S}^T \mathbf{U}^T \mathbf{U} \mathbf{S} \mathbf{V}^T \\ &= \mathbf{V} \mathbf{S}^T \mathbf{S} \mathbf{V}^T \\ &= \mathbf{V} \mathbf{D} \mathbf{V}^T \end{aligned}$$

$$(\mathbf{X}^T \mathbf{X}) \mathbf{V} = \mathbf{V} \mathbf{D}$$



PCA algorithm

1. Normalise the data to be zero-mean.
2. Calculate the sample covariance matrix.
3. Find the D eigenvector-eigenvalue pairs of the sample covariance matrix.
4. Choose the M eigenvectors corresponding to the highest eigenvalues.
5. Project the data to the lower-dimensional space.

Videos covered in this note

- [PCA 1 - Introduction](#) (16 min)
- [PCA 2 - Mathematical background](#) (7 min)
- [PCA 3 - Setup](#) (17 min)
- [PCA 4 - Learning](#) (19 min)
- [PCA 5 - Minimising reconstruction](#) (7 min)
- [PCA 6 - Relationship to SVD](#) (9 min) (not examinable)
- [PCA 7 - Steps](#) (6 min)

Reading

- ISLR 12.2 (excluding 12.2.3, although this is interesting)